

Transit infrared spectroscopy of the hot Neptune around GJ 436 with the *Hubble Space Telescope*★

F. Pont,^{1†} R. L. Gilliland,² H. Knutson,³ M. Holman³ and D. Charbonneau³

¹University of Exeter, The Queens Drive, Exeter, Devon, EX4 4QJ

²Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

³Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

Accepted 2008 October 29. Received 2008 October 29; in original form 2008 August 18

ABSTRACT

The nearby transiting system GJ 436b offers a unique opportunity to probe the structure and atmosphere of an extrasolar ‘hot Neptune’. In this Letter, we present the main results of observations covering two transit events with the Near Infrared Camera and Multi Object Spectrograph (NICMOS) camera on the *Hubble Space Telescope* (*HST*). The data consist in high-cadence time series of grism spectra covering the 1.1–1.9 μm spectral range. We find $R_{\text{pl}} = 4.04 \pm 0.10 R_{\oplus}$ and $R_{\star} = 0.446 \pm 0.011 R_{\odot}$ for the planet and star radius, confirming and improving earlier measurements with a ground-based photometry and a *Spitzer* light curve at 8 μm , as opposed to a much higher value obtained with the Fine Guidance Sensor on the *HST*. We measure no departure from strict periodicity in the transits to the level of ~ 7 s. This strongly disfavours the proposed explanation of the orbital eccentricity of GJ 436b in terms of the perturbation by another close-by planet. We measure a flat transmission spectrum at the level of a few parts per 10 000 in flux, with no significant signal in the 1.4- μm water band to a level comparable to the maximum amplitude of the effect predicted by planetary atmosphere models.

Key word: planetary systems.

1 INTRODUCTION

The planet orbiting around the nearby M-dwarf star GJ 436 is the first known transiting Neptune-class planet – and to date the only one (Butler et al. 2004; Gillon et al. 2007; Maness et al. 2007). The planet has a 2.64 d orbit with an eccentric of 0.15, and a mass of ~ 23 Earth masses. Its radius of $\sim 4R_{\oplus}$ implies a primarily icy or rocky composition with an envelope of hydrogen and helium (Adams, Seager & Elkins-Tanton 2008), a hotter version of Uranus or Neptune. Measurements of the secondary eclipse in the mid-infrared with the *Spitzer Space Telescope* indicate a photospheric temperature in the vicinity of 700 K (Deming et al. 2007).

A second planet has been hypothesized to explain the high orbital eccentricity (e.g. Ribas, Font-Ribera & Beaulieu 2008a). This planet would induce variations in the transit timing detectable with sufficiently precise transit photometry.

Spectroscopic time series during transit can also provide information on the transmission spectrum of the planetary atmosphere: in wavelengths where its atmosphere is more opaque, the planet appears larger across the stellar disc and the transit is slightly deeper.

This effect was first detected for the transiting system HD 209458 by Charbonneau et al. (2002), and recently used to obtain an extensive characterization of the atmospheric transmission spectrum of HD 189733b (Pont et al. 2007; Swain, Vasisht & Tinetti 2008).

We here present the analysis of two sequences of spectra obtained with the Near Infrared Camera and Multi Object Spectrograph (NICMOS) camera on the *Hubble Space Telescope* (*HST*) with the G141 grism, during two transits of the GJ 436 system in 2007 November 11 and December 15. These data pertain to the issues mentioned above: the resulting precise time series in total flux provides very accurate constraints on the stellar and planetary radius, and determine the timing of the two transits to a few seconds, the spectroscopic time series constrains the planetary atmosphere transmission spectrum.

2 DATA AND REDUCTION

The images were taken with the G141 grism on the NICMOS camera, sampling the 1.1–1.9 μm spectral interval. The first *HST* visit consists of 937 images taken between JD = 2454415.48 and 2454415.71, the second visit of 917 images between 2454449.85 and 2454450.08. The images are 1.993-s exposures obtained on 11-s centres. Since GJ 436 is not situated in the continuous viewing zone of *HST*, data were acquired during only half of the orbits. The timing of the observations is such that orbit 3 (of 4) of each visit occurs during the transit, sampling the transit centre and

★Based on observations with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS 5-26555.

†E-mail: fpont@astro.ex.ac.uk

egress. Orbit 2 and 4 are used to set the out-of-transit baseline and calibrate the instrumental effects. As in previous similar observations, orbit 1, during which the telescope and instrument settle to the new mode, cannot be used at this level of accuracy. We used the same defocus setting as in previous NICMOS grism observations of bright transiting planets. The highest signal peak on the detector is 87 000 electrons, which corresponds to 43 per cent of the 215 000 full well-depth for the NIC3 detector.

2.1 Flat-fielding and flux extraction

The standard pipeline was used to perform bias level, zero read subtraction, non-linearity correction, dark subtraction and cosmic ray removal. Wavelength-dependent flat-field corrections were computed by interpolating between a set of calibration flat-fields taken in different filters.

To calculate the flux, we define an extraction box out to a specified intensity level chosen by running several trials and selecting the one that returns the smallest time series rms over out-of-transit orbits 2 and 4 of each visit. We interpolate spatially over a small set of dead and extra noisy pixels. A global sky level is then evaluated and subtracted for each image.

2.2 External parameter decorrelation

Even though the instrumental setup, pointing and observing conditions were kept very stable, the photon signal-to-noise ratio is so high in our data set that it is important to correct even tiny instrumental effects.

As in Pont et al. (2007), we removed these effects to first order with multilinear decorrelation. We track four external variables for each exposure: the drift of the spectrum in x and y coordinates on the detector, its rotation and width change. The shift in x -position was evaluated by cross-correlating extracted 1-d spectra. The changes in y -position, width and rotation were defined by fitting a set of 1-d Gaussian in the cross-dispersion direction along the first-order spectrum. The amplitude of the changes was ± 0.04 pixel in position, ± 0.03 pixel in width and ± 0.05 degrees in rotation in the first visit, and about twice as large in the second visit. To these four vectors, we add a linear time vector as a fifth decorrelation parameter (to account for a constant drift over the whole observing sequence). For each of the two visits, we calibrate the corrections from a multilinear regression of the flux residuals against the value of the external parameters for the second and fourth orbit (the first orbit is discarded and the third consists of in-transit data).

This external parameter decorrelation decreases the rms of the intensity in 1-min time intervals from 3.27×10^{-4} to 1.97×10^{-4} for the first visit and 5.68×10^{-4} to 2.07×10^{-4} for the second. Contrarily to the case of ACS data on HD 209458 (Pont et al. 2007), the latter values are still two to three times larger than the photon-noise limit (0.74×10^{-4}).

Nevertheless, the decorrelation produces satisfactory results for the data of the first visit, as shown a posteriori by the residuals around a transit light-curve fit. The results are less well behaved for the second visit. The reason is that several external parameters take values during the third orbit (the in-transit orbit), that are not covered during the second and fourth orbits used for decorrelation. Therefore, the decorrelation is performed by extrapolating the dependence between parameters and residuals outside the range actually sampled.

2.3 Correlated noise analysis

Noise on transit time series has a very different impact depending on its covariance properties, as discussed in Pont, Zucker & Queloz (2006). Noise correlated in time and frequency, such as that produced by telescope and detector systematics, produces much higher uncertainties on the final results than uncorrelated noise sources such as the photon shot noise, because correlated errors do not average out quickly with higher number of exposures.

We model the noise as a combination of purely random and entirely correlated components, following the approach of Pont et al. (2006):

$$\sigma_{\text{tot}}^2 = \sigma_w^2 + \sigma_t^2 + \sigma_v^2$$

with σ_w the purely white noise component, σ_t and σ_v the components correlated in time and frequency, respectively. The coefficients are estimated by ensuring $\sigma^2 = (\sigma_w^2 + \sigma_v^2)/N + \sigma_t^2$ in 15-min time bins and $\sigma^2 = (\sigma_w^2 + \sigma_t^2)/N + \sigma_v^2$ in 14-pixel bands along the dispersion direction (14 pixel is the approximate point spread function size). We find $\sigma_{\text{tot}} = 4.7 \times 10^{-4}$, the total rms of individual intensity values, breaks down into $\sigma_w = 3.9 \times 10^{-4}$, $\sigma_t = 2.3 \times 10^{-4}$ and $\sigma_v = 1.0 \times 10^{-4}$. The white-noise term averages out when integrated over large number of exposures. For global parameters like the planet-to-star radius ratio, the second terms (noise with time correlation) will dominate the error budget. We thus expect the effect of correlated noise to amount to $\sim 3 \times 10^{-4}$ on the transit depth, corresponding to a precision of 0.6 per cent on the radius ratio of the system. For the transmission spectrum, the most relevant source of noise is the last one (noise with wavelength correlation). The noise budget is $\sim 1.5 \times 10^{-4}$ on transmission spectral features including white and correlated noise.

3 INTEGRATED-LIGHT LIGHT-CURVE ANALYSIS

3.1 Transit parameters

The light curve integrated over the whole spectrum is plotted in Fig. 1. The residuals have the same amplitude and correlation properties inside and outside the transit, indicating no significant deviations from the signal expected for the transit of an opaque body in front of an M-dwarf star. In particular, no indication of the transit of another, smaller body in front of the star is seen, nor in-transit residuals that could indicate that the planet is occulting a large star spot (as was the case for HD 189733 in Pont et al. 2007).

Since our data set covers the transit egress twice and never the ingress, it is not suited to measure the position of the mean epoch of the transit. We therefore adopt the epoch and period of the transit signal obtained on five partial transits with the Fine Guidance Sensor of the *HST* by Bean et al. (2008; hereafter B08): $P = 2.6438904$ d and $T = 2454455.27924$ HJD. We also adopt the eccentricity and argument of periastron from the radial-velocity orbit. At the level of accuracy reached by our data, it is important to use a precise limb-darkening profile, even though the effect of limb darkening is somewhat lower in the near-infrared (near-IR) than at shorter wavelengths. To calculate the limb-darkening parameters, we used a model selected from a grid of Kurucz atmosphere models with an effective temperature of 3500 K, $\log(g) = 5.0$, a turbulent velocity of 1 km s^{-1} and $[\text{Fe}/\text{H}] = 0.0$. We calculate limb darkening numerically on 17 points along a chord crossing the stellar disc, and fit the four-coefficient non-linear law of Claret (2000). We calculate the limb-darkening coefficients for the central wavelengths of 114 one-pixel




Figure 1. NICMOS integrated-light time series (decorrelated for instrumental effects), in 1-min bins (five to six data points per bin), with best-fitting transit model light curve, for the two *HST* visits. The rms around the curve is 1.97×10^{-4} for the first visit and 2.07×10^{-4} for the second.

intervals along the spectral dispersion of the NICMOS spectra, then compute the intensity-weighted mean of the coefficients for the integrated-light light curve. We find (1.533, -2.234 , 1.913 and -0.643) for the four coefficients.

We fit a transit curve model using the Mandel & Agol (2002) algorithm on the NICMOS light curve, obtaining $R_{\text{pl}}/R_* = 0.0831 \pm 0.0005$, $T_{\text{tr}} = 0.0317 \pm 0.0004$ d and $b = 0.850 \pm 0.007$ for the geometric transit parameters (radius ratio, transit duration¹ and impact parameter). The uncertainties are calculated using the method of Pont et al. (2006), accounting for the presence of correlated noise in the data. Fitting the two transits separately yields $R_{\text{pl}}/R_* = 0.0832 \pm 0.0006$, $T_{\text{tr}} = 0.0317 \pm 0.0004$ and $b = 0.857 \pm 0.013$ for the first visit and $R_{\text{pl}}/R_* = 0.0830 \pm 0.0006$, $T_{\text{tr}} = 0.0320 \pm 0.0004$ and $b = 0.843 \pm 0.014$ for the second visit. The two sets of values are compatible within their error bars.

We can then iterate on the transit epoch and period to attain a self-consistent solution. With the geometric transit parameters fixed, we free only the two transit timings for the NICMOS visits, findings: $T_1 = 2454415.62074 \pm 0.00008$ and $T_2 = 2454449.99141 \pm 0.00008$. These values are not significantly different from those obtained by extrapolating from the epoch and period found by B08 (5 ± 7 s earlier and 2 ± 7 s later) and used in the first fit. Further iterating is thus unnecessary.

3.2 Transit timing

Fig. 2 shows the transit central times of the NICMOS and fine guidance sensor (FGS) data, as residuals compared to the solution of B08, that also take into account the earlier *Spitzer* and ground-based measurements. The transit signal is found to be precisely periodic to the level of a few seconds. Such stability severely constrains




Figure 2. Transit timings for our two transits (smaller error bars) and the five transits with FGS from B08. The dotted line shows the expectation for a strictly periodic signal compatible with the earlier *Spitzer* transit.

the presence of a second massive planet with a period comparable to that of GJ 436b, as invoked, for instance, by Ribas et al. (2008a) to account for radial-velocity residuals and the orbital eccentricity. This scenario (and subsequent variations to account for new observations, e.g. Ribas et al. 2008b) predicts oscillations of several minutes in the transit times, which, barring highly unlikely coincidences, are excluded by the NICMOS data.

We performed numerical integrations similar to those in Agol et al. (2005) and Holman & Murray (2005) to estimate the amplitude of transit timing variations that would be induced by an additional planet in the GJ 436 system. A perturbing planet with mass $0.01 M_{\oplus}$ in either the 2:1 or 3:1 mean motion resonance will generate rms transit timing variations of ~ 7 s, assuming that the perturbing planet shares the orbital plane, argument of pericenter and eccentricity ($e=0.15$) of GJ 436b. Likewise, a $5 M_{\oplus}$ planet in a similar but non-resonant orbit between the 2:1 and 3:1 mean motion resonance will generate ~ 7 s rms timing variations.

Bean & Seifahrt (2008) discuss the possible location in the $M_p - a$ plane of a second planet accounting for the eccentricity of GJ 436b (see their fig. 1). Our data, while not strictly excluding all scenarios, further narrow down the parameter space available. If GJ 436b's unusual eccentricity is not the result of perturbations from a second planet in the system, we must consider the possibility that the planet's eccentricity is primordial and we have simply underestimated the circularization time-scale for this system. This would require an unusually high value for the planet's tidal Q factor, but this factor is generally poorly constrained for extrasolar planets (Ogilvie & Lin 2004; Wu 2005).

3.3 Planetary radius

Adopting $M_* = 0.452 \pm 0.013$ from Torres (2007), the transit parameters found from the NICMOS data imply $R_* = 0.446 \pm 0.011 R_{\odot}$ for the stellar radius and $R_{\text{pl}} = 4.04 \pm 0.10 R_{\oplus}$ (25750 ± 650 km) for the planetary radius. The systematic uncertainties due to stellar evolution models (for M_*) are not included in the error estimates.

Our value for the planetary radius is in agreement with initial ground-based estimates and the value derived from the *Spitzer* coverage at $8 \mu\text{m}$ ($R_{\text{pl}} \simeq 4.0\text{--}4.3 R_{\oplus}$). We do not confirm the significantly higher value obtained by B08 ($R_{\text{pl}} = 4.90^{+0.45}_{-0.33} R_{\oplus}$) from the FGS light curves. B08 discuss several possible explanations for either the *Spitzer* or the FGS radius to be off by $\sim 2\sigma$. In particular, the FGS light curve is built from five different partial transits, with relative shifts between the different transits as free parameters. Unrecognized time-dependent effects (instrumental systematics

¹ Transit time of the centre of the planet across the stellar disc.

as well as stellar variability and spots) could thus modify the shape of the transit. The effect of stellar limb darkening is also much stronger in the visible wavelengths detected by FGS than in the near-IR, and more difficult to calculate from models – especially for an M-type spectrum. The effect of stellar variability and spots is also larger for shorter wavelengths. Given these factors and the higher total signal-to-noise ratio of the NICMOS data, the radius value that we find suggests that the FGS radius is indeed overestimated.

4 SPECTROSCOPIC ANALYSIS

The primary objective of our observations was to provide broad constraints on the transmission spectrum of the planetary atmosphere in the 1.1–1.9 μm range, in particular around the water bands near 1.4 μm .

Models for this planet assuming a primarily hydrogen-dominated atmosphere with solar metallicity predict that water will exist in gas phase in the upper atmosphere, as this region is too hot for water vapour clouds and too cool for water to be thermally dissociated. At the predicted abundances, this water vapour produces a strong absorption signal at near-IR wavelengths, including one band at 1.4 μm that falls in the centre of our spectrum.

We can estimate in an approximate way the size of the expected absorption signal by calculating the atmospheric scaleheight, which is given by $H = kT/(g\mu)$, where T is the temperature, g is the surface gravity and μ is the mean molecular weight of the atmosphere. For GJ 436b, assuming a temperature of 710 K (Deming et al. 2007; Demory, Gillon & Barman 2007), a surface gravity of 1280 cm s^{-1} (Butler et al. 2004; Torres 2007) and an atmosphere of molecular hydrogen, this would correspond to a scaleheight of 230 km. The change in the depth of the transit is proportional to the additional area occulted, typically approximated as $10 \cdot H$ (Seager & Sasselov 2000), so the total size of the signal would be 2×10^{-4} . This is consistent with the predictions from full 1D radiative transfer atmosphere models for GJ 436b (Miller-Ricci & Sasselov, private communication), which indicate that the predicted signal from water absorption in our selected bandpass should be $0.6\text{--}1.1 \times 10^{-4}$, where the lower end of this range corresponds to an atmosphere with 30 times solar metallicity and the upper end is for an atmosphere with solar metallicity (see Miller-Ricci, Seager & Sasselov 2008 for a full description of the methodology used for these models). Uniformly increasing the fraction of heavy elements in this planet's atmosphere does not significantly alter the relative depths of the various absorption features, but it does produce a net decrease in the strength of those features as the increased value of μ reduces the atmospheric scaleheight.

To recover the wavelength-dependent information from the NICMOS data, we repeated the extraction along columns on the CCD perpendicular to the direction of dispersion. We built a differential indicator of the intensity of absorption in the 1.4- μm water band, by computing the difference between the flux in a 1.34–1.53 μm passband with the mean of the flux in two side bands, 1.21–1.30 and 1.55–1.64 μm . We correct for the different limb-darkening coefficients in these passbands with the Kurucz models.

The presence of a planetary atmosphere with excess opacity in the water band would leave a transit-like signature in the run of this indicator with time, because the planet would appear larger within the wavelength of the band than outside. We find a flux *excess* of $+1.42 \pm 0.89 \times 10^{-4}$ in the water passband during the transit. The uncertainty is estimated from the variance of the time series of the water indicator, and does not include the uncertainties in the cor-

rections of instrumental systematics. Based on the noise analysis of Section 2.3, we estimate these at $\sim 1.0 \times 10^{-4}$. To sum up, we find a $\sim 1\sigma$ signal in the direction opposite to the expected water absorption band, with a comparable amplitude. Thus, although spectral signatures much larger than that predicted by the models are ruled out, the NICMOS data are not sufficient to measure spectral features of the planetary atmosphere.

We also built a low-resolution transit spectrum from our data, to check whether any feature in the planet transmission spectrum was visible above the uncertainties introduced by residual instrumental systematics. We binned the column-by-column wavelength extraction in 5-pixel bands (0.04- μm passbands), and fitted a transit shape with all parameters fixed to the best-fitting values and theoretical limb-darkening coefficients, except the transit depth. Wavelength-dependent systematics in the time series were corrected for each passband to first order with a five-parameter description: two quadratic functions of *HST* orbital phase before and after the moment when the telescope enters the Earth's shadow and a zero-point drift. We retain only the data when the covariance between the decorrelation parameters and the transit depth was small, restricting the results to the $1.35 < \lambda < 1.85 \mu\text{m}$ interval and the first *HST* visit. Outside this wavelength interval, the resulting spectrum depends significantly on the parameters used in the decorrelation. For the second visit, the wavelength-dependent decorrelation is not stable, because the pointing and focus of the telescope took values during the in-transit orbit that are incompletely sampled by the orbits before and after the transit.

Fig. 3 plots the results in terms of transit cross-section ($R_{\text{pl}}^2/R_{\text{star}}^2$), together with the results from ground-based observations (Shporer et al. 2008), the *Spitzer* 8.0- μm measurement (Southworth 2008),




Figure 3. Planet cross-section (in percentage of the stellar disc covered) as a function of wavelength for GJ 436b. The NICMOS data (squares) is binned over 5-pixel intervals across the spectral dispersion direction. The short-wavelength points come from a collection of ground-based data (Shporer et al. 2008, triangle) and the B08 data (refitted with our orbital parameters). The long-wavelength point is the 8.0- μm *Spitzer* transit (Southworth 2008). Error bars for space data do not include the uncertainties on the correction of instrumental systematics.

Table 1. New and updated parameters of the GJ 436 system.

Period (d)	2.6438904 (from B08)
Radius ratio	0.0831 ± 0.0005
Impact parameter	0.850 ± 0.007
Transit duration (d)	0.0317 ± 0.0004
Transit timing (HJD)	$2454415.62074 \pm 0.00008$
	$2454449.99141 \pm 0.00008$
Star radius (R_{\odot})	0.446 ± 0.011
Planet radius (R_{\oplus})	4.04 ± 0.10

and the B08 result refitted with our transit parameters. The dotted line shows the best-fitting cross-section from the integrated-light NICMOS data. The NICMOS spectrum has a rms of 1.9×10^{-4} , compared to an expected noise of 1.4×10^{-4} (including 1.0×10^{-4} noise correlated in wavelength). The data can be explained by a constant radius over all the wavelength range, and there is no compelling evidence at this point for spectral features that could be attributed to the effect of the planetary atmosphere at the level of a few parts per 10^4 .

5 CONCLUSION

Two transits of the hot Neptune GJ 436b were measured in very high signal-to-noise ratio grism spectroscopy in the 1.1–1.9 μm range with NICMOS on the *HST*. The transit shape indicates a size of $R_{\text{pl}} = 4.04 \pm 0.10 R_{\oplus}$ for the planet (assuming $M_* = 0.452 \pm 0.013 M_{\odot}$), confirming the values obtained with the *Spitzer* 8- μm data, rather than the higher radius from FGS/*HST* data in the visible. The higher FGS value is probably due to the instrumental systematics, unexpected values of limb darkening or irregularities on the stellar surface.

No significant departures from a pure, strictly periodic transit signal are present beyond the level of time-correlated residuals from instrumental systematics ($\sim 10^{-4}$ in flux), indicating the absence of strong intensity fluctuations on the surface of the star crossed by the planet, and of transit timing variations on the scale of a few dozen seconds. This is a strong argument against a second close-in planet producing the perturbations responsible for the eccentricity of the orbit of GJ 436b.

We analyse the wavelength dependence of the transit depth in search of features introduced by the planetary atmosphere. We find no such feature at the level of a few parts per 10^4 , and measure a flux excess of $+1.42 \pm 0.89 \times 10^{-4}$ in the water vapour absorption band around 1.4 μm (where models predict a signal of 10^{-4} or lower in the opposite direction). Therefore, the data exclude an anomalously large signal in transmission spectroscopy in this band, but are not sufficient to measure the actual size of the water absorption feature. Our ability to remove systematics from the data is ultimately the limiting factor in this analysis; if we are to achieve a detection of water vapour in this planet's atmosphere we will have to improve our corrections for these effects in future analyses of *HST* observations and reach a noise level much closer to that of the photon-noise-limited precision for this system.

ACKNOWLEDGMENTS

The authors wish to thank Eliza Miller-Ricci and Sara Seager for atmosphere models provided in advance of publication and insightful comments, Michel Mayor for support with the observing proposal, Daniel Fabrycky for helpful discussions about transit timing variations in the GJ 436 system and the anonymous referee for numerous helpful comments.

REFERENCES

- Adams E. R., Seager S., Elkins-Tanton L., 2008, *ApJ*, 673, 1160
 Agol E., Steffen J., Sari R., Clarkson W., 2005, *MNRAS*, 359, 567
 Bean J. L., Seifahrt A., 2008, *A&A*, 487, L25
 Bean J. L. et al., 2008, *A&A*, 486, 1039 (B08)
 Butler R. P., Vogt S. S., Marcy G. W., Fischer D. A., Wright J. T., Henry G. W., Laughlin G., Lissauer J. J., 2004, *ApJ*, 617, 580
 Charbonneau D., Brown T. M., Noyes R. W., Gilliland R. L., 2002, *ApJ*, 568, 377
 Claret A., 2000, *A&A*, 363, 1081
 Deming D., Harrington J., Laughlin G., Seager S., Navarro S. B., Bowman W. C., Horning K., 2007, *ApJ*, 667, 199
 Demory B.-O., Gillon M., Barman T., 2007, *A&A*, 475, 1125
 Gillon M. et al., 2007, *A&A*, 472, L13
 Holman M. J., Murray N. W., 2005, *Sci*, 307, 1288
 Mandel K., Agol E., 2002, *ApJ*, 580, 171
 Maness H. L., Marcy G. W., Ford E. B., Hauschildt P. H., Shreve A. T., Basri G. B., Butler R. P., Vogt S. S., 2007, *PASP*, 119, 90
 Miller-Ricci E., Seager S., Sasselov D., 2008, *ApJ*, in press (arXiv:0808.1902)
 Ogilvie G. I., Lin D. N. C., 2004, *ApJ*, 610, 477
 Pont F., Zucker S., Queloz D., 2006, *MNRAS*, 373, 231
 Pont F. et al., 2007, *MNRAS*, 385, 109
 Ribas I., Font-Ribera A., Beaulieu J.-P., 2008a, *ApJ*, 677, 59
 Ribas I., Font-Ribera A., Beaulieu J.-P., Morales J. C., Garcia-Melendo E., 2008b, in Pont F., Sasselov D., Holman M., eds, *Proc. IAU Symp.* 253, in press
 Seager S., Sasselov D. D., 2000, *ApJ*, 537, 9
 Shporer A., Mazeh T., Winn J. N., Holman M. J., Latham D. W., Pont F., Esquerdo G. A., 2008, *A&A*, preprint (arXiv:0805.3915)
 Southworth J., 2008, *MNRAS*, 386, 1644
 Swain M., Vasisht G., Tinetti G., 2008, *Nat*, 452, 329
 Tinetti G. et al., 2007, *Nat*, 448, 169
 Torres G., 2007, *ApJ*, 671, 65
 Wu Y., 2005, *ApJ*, 635, 674

This paper has been typeset from a \LaTeX file prepared by the author.